

A large local rotational speed for the Galaxy found from proper-motions: Implications for the mass of the Milky-Way

René A. Méndez

Cerro Tololo Inter-American Observatory, Casilla 603, La Serena, Chile.

E-mail: rmendez@noao.edu

Imants Platais¹, Terrence M. Girard, Vera Kozhurina-Platais, William F. van Altena

Department of Astronomy, Yale University, P. O. Box 208101, New Haven, CT 06520-8101.

E-mail: imants, vera, girard, vanalten @astro.yale.edu

ABSTRACT

Predictions from a Galactic Structure and Kinematic model are compared to the absolute proper-motions of about 30,000 randomly selected stars with $9 < B_J \leq 19$ derived from the Southern Proper-Motion Program (SPM) toward the South Galactic Pole. The absolute nature of the SPM proper-motions allow us to measure not only the relative motion of the Sun with respect to the local disk, but also, and most importantly, the overall state of rotation of the local disk with respect to galaxies. The SPM data are best fit by models having a solar peculiar motion of $+5 \text{ km s}^{-1}$ in the V-component (pointing in the direction of Galactic rotation), a large LSR speed of 270 km s^{-1} , and a disk velocity ellipsoid that points towards the Galactic center. We stress, however, that these results rest crucially on the assumptions of both axisymmetry and equilibrium dynamics.

The absolute proper-motions in the U-component indicate a solar peculiar motion of $11.0 \pm 1.5 \text{ km s}^{-1}$, with no need for a local expansion or contraction term.

The implications of the large LSR speed are discussed in terms of gravitational mass of the Galaxy inferred from the most recent and accurate determination for the proper-motion of the LMC. We find that our derived value for the LSR is consistent both with the mass of the Galaxy inferred from the motion of the Clouds ($3 - 4 \times 10^{12} M_\odot$ to $\sim 50 \text{ kpc}$), as well as the timing argument, based on the binary motion of M31 and the Milky Way, and Leo I and the Milky Way ($\geq 1.2 \times 10^{12} M_\odot$ to $\sim 200 \text{ kpc}$).

Subject headings: Astrometry: stellar dynamics – stars: kinematics – stars: fundamental parameters – Galaxy: fundamental parameters – Galaxy: kinematics and dynamics – Galaxy: structure

¹Visiting Astronomer, Cerro Tololo Inter-American Observatory. CTIO is operated by AURA, Inc. under contract to the National Science Foundation.

1. Introduction

In this paper, we present the main results from an analysis of the kinematic data obtained in the context of a massive *absolute* proper-motion survey, the Southern Proper-Motion Program (SPM hereafter). Our analysis is based on the SPM Catalog as described by Platais et al. (1998). This catalog provides positions, absolute proper-motions, and *BV* photometry for about 30,000 randomly selected stars, among other objects, near the South Galactic Pole. The sky coverage of the SPM Catalog is about 720 deg^2 in the magnitude range $5 < V < 18.5$. The accuracy of individual absolute proper-motions is $3\text{--}8 \text{ mas yr}^{-1}$ depending on the star's magnitude. For a complete description of the catalogue structure, contents, plate measurement and other astrometric/photometric details, the reader is referred to Girard et al. (1998, Paper I) and Platais et al. (1998, Paper II). A detailed analysis of the velocity distribution function, and star- and color-counts for stars from the disk, thick-disk, and halo derived from the SPM photometry and astrometry is addressed by Méndez et al. (1999, Paper III).

The arrangement of this Letter is the following: In Section 2 we present an overview of the SPM photometric and proper-motion data. In Section 3 the model used to compare the observed proper-motion distributions with those predicted is described, while Section 4 presents the model predictions as compared to the proper-motion data. Section 5 presents the implications of our results for the global mass of the Galaxy. Finally, Section 6 presents the main conclusions of the paper.

2. The randomly selected sample

The random SGP-SPM sample has been chosen such that, at a given B_J magnitude interval, a fixed number of stars are randomly extracted from all stars available in that magnitude interval. As a result, one-magnitude intervals are chosen in the range $9 \leq B_J \leq 19$ for the subsequent analysis. At a given magnitude interval, the complete sample and the randomly selected sample differ only by a scale factor. Therefore, color and kinematic properties binned in the proper magnitude intervals should be very similar to those derived from a complete sample, except for the increased uncertainties on the derived values because of the smaller sample - an effect that is fully taken into account in the analysis below.

3. The starcounts and kinematic model

We model the starcounts concurrently with the kinematics by using the Galactic structure and kinematic model presented by Méndez and van Altena (1996). The starcount model employed here has been tested under many different circumstances, and has proved to be able to predict starcounts that match the observed magnitude and color counts (in both shape *and* number) to better than 10%, and in many cases to better than 1%. The kinematic model presented by Méndez and van Altena (1996) has also been shown to be able to reproduce the kinematics of disk stars toward different Galactic fields. In Méndez et al. (1999), a full analysis of the SPM-SGP data in terms of star and color-counts, as well as the kinematics is presented. A basic assumption in the Méndez and van Altena (1996) model is that of the Gaussian shape of the velocity ellipsoid. While the velocity distribution observed for late-type stars (e.g., Dehnen 1998) is clearly skewed and cannot be properly modeled by a Gaussian, the error in the predicted median proper motion made by using a shifted Gaussian is likely to be small. Indeed, Méndez et al. (1999), show that shifted-Gaussian velocity distribution functions do provide a good fit to the full SPM proper-motion distributions as a function of apparent magnitude, with perhaps a slight predicted excess of stars with positive proper-motions in the direction of Galactic rotation.

Because of projection effects near the SGP, our model comparisons are most sensitive to the motions in the direction of Galactic rotation (the so-called V-component), and along the Galactic center-anticenter direction (the U-component), and *not* to the motions perpendicular to the Galactic plane (the W-component).

3.1. The Solar peculiar velocity and the motion of the LSR

There have been a number of determinations for the solar peculiar motion. The classical result, quoted in Mihalas and Binney (1981), gives $(U_\odot, V_\odot, W_\odot) = (+9.0, +12.0, +7.0) \text{ km s}^{-1}$, essentially based upon Delhaye's (1965) compilation. More recent values for the solar peculiar motion do not seem to have converged to a single value, especially in the V-component. Dehnen and Binney (1998) find $(U_\odot, V_\odot, W_\odot) = (+10.00 \pm 0.36, +5.25 \pm 0.62, +7.17 \pm 0.38) \text{ km s}^{-1}$ from a carefully selected unbiased sample of Hip-

parcos stars, while Miyamoto and Zhu (1998) find $(U_{\odot}, V_{\odot}, W_{\odot}) = (+10.62 \pm 0.49, +16.06 \pm 1.14, +8.60 \pm 1.02)$ km s⁻¹ from 159 Cepheids, *also* from the Hipparcos catalogue. As suggested earlier by Robin and Oblak (1987), such variation in the $V_{\odot}$ component is probably due to difficulties in separating the asymmetric drift from the intrinsic solar motion, and also because of the peculiar motions exhibited by the very young OB stars, which are still moving under the influence of the spiral arm kinematics and/or of their parent molecular cloud.

In 1985, the IAU adopted a value for the motion of the LSR of 220 km s⁻¹. Kerr and Lynden-Bell (1986) have discussed extensively the determinations of $V(R_0)$, as well as R_0 , and the Oort constants A and B available until then. From a straight mean of different determinations they obtained $V_{\text{LSR}}(R_0) = 222 \pm 20$ km s⁻¹ (their Table 4), while from independent determinations of R_0 , A, and B (their Tables 3 and 5), we find $V_{\text{LSR}}(R_0) = 226 \pm 44$ km s⁻¹. Although the uncertainties involved in $V_{\text{LSR}}(R_0)$ are larger than those of $V_{\odot}$, we shall see that the relative motion of disk stars is more affected by uncertainties in the Solar peculiar motion than uncertainties in the motion of the LSR, since the whole nearby disk population is approximately co-moving with the LSR.

4. Model comparisons to the absolute proper-motions

The basic standard kinematic parameters employed in the model are described in Méndez and van Altena (1996) and Méndez et al. (1999). The standard model adopts $(U_{\odot}, V_{\odot}, W_{\odot}) = (+11.0, +14.0, +7.5)$ km s⁻¹ for the peculiar solar motion, and a flat rotation curve with $V_{\text{LSR}}(R_0) = +220$ km s⁻¹. Because we are analyzing data near the SGP, our model predictions do not span a large range in Galactocentric distance, and therefore the model is insensitive to the value adopted for the *slope* of the rotation curve near R_0 (although see Méndez et al. 1999). As for the disk kinematics, we have adopted the velocity dispersions given in Méndez and van Altena (1996), with the scale-heights adopted in Méndez et al. (1999) which reproduce the observed SPM color- and magnitude counts, a scale-length of 3.5 kpc (again, the model is not sensitive to this last parameter, as it enters as a function of the Galactocentric distance, which for the SPM-SGP data is quasi-constant), and a value of $q = 0$ for the velocity ellipsoid (see Eq. 4 in Méndez and van Altena

1996); $q = 0$ implies a velocity ellipsoid parallel to the Galactic plane at all heights from the Galactic plane, while $q = 1$ is for a velocity ellipsoid that points towards the Galactic center). As shown by Méndez et al. (1999) the exact choice of kinematical parameters for the thick-disk and halo are not critical to the conclusions of this Letter. The mean velocity of stars with different velocity dispersion is computed via the proper moments of the Collisionless Boltzmann equation (CBE). The CBE allow us to relate our adopted velocity dispersions to the density laws, and to derive an expression for the velocity lag as a function of the net rotation of the disk (the local LSR) and the pressure support provided by the velocity dispersions (see, e.g., Kuijken and Gilmore 1989, Gilmore et al. 1989). We must emphasize however that, if the data contain information that is inconsistent with the assumptions made (in particular, the validity of an axisymmetric CBE, and the neglect of stellar streams), then these assumptions inevitably could lead to biased results – the classical problem of all parametric methods.

4.1. The Solar motion and the LSR speed

Figures (1) and (2) show the model-predicted median absolute proper-motions for the standard model parameters compared to the observations. These figures clearly show that, while the standard model produces a very good fit to the U-component of the proper-motion (along the Galactocentric direction), the V-component (along Galactic rotation) is grossly underestimated, especially for $B_J < 14$, where the disk component dominates the overall kinematics.

A radical change in the model predictions occurs when we change the peculiar Solar motion from the standard value of $V_{\odot} = +14$ km s⁻¹ to $V_{\odot} = +5$ km s⁻¹. The largest effect occurs at the brightest bins, i.e., for nearby disk stars where the Solar peculiar motion dominates the reflex motion. On the other hand, we may keep the conventional Solar peculiar velocity, but change the overall rotation speed for the LSR from the IAU adopted value of $+220$ km s⁻¹ to $+270$ km s⁻¹, as suggested by recent Hipparcos results (Miyamoto and Zhu 1998). In this case, the change in the predicted motion becomes more important at fainter magnitudes where one is sampling objects from the other Galactic components located at larger distances, and where the dominant effect is that of the overall rotation of the disk with respect to galaxies, and the relative state of rotation between the different Galactic components. At the brightest

bins, the effect of changing $V_{\text{LSR}}(R_0)$ becomes also noticeable because of the larger fraction of bright giants, which can be seen to large distances and, hence, the differential rotation effects become amplified. A model in which we simultaneously change the Solar peculiar motion *and* the LSR rotational speed to $+5 \text{ km s}^{-1}$ and $+270 \text{ km s}^{-1}$, respectively, produces a much better fit to the observed motions. We note that, in this model, the velocity lags for the thick-disk and halo are increased in proportion to the increase of the disk's rotational speed, as the net rotation of those two components is kept constant at 180 km s^{-1} and 0 km s^{-1} , respectively (for details see Méndez et al. 1999). We conclude that a model with $V_{\odot} = +5 \text{ km s}^{-1}$ and $V_{\text{LSR}}(R_0) = 270 \text{ km s}^{-1}$ clearly provides a much better fit to the median motion in V than does the standard model with $V_{\odot} = +11 \text{ km s}^{-1}$ and $V_{\text{LSR}}(R_0) = 220 \text{ km s}^{-1}$.

The predicted median proper-motion in the U-component shows a good fit to the observed values, and the changes in the V-component described above do not affect this parameter substantially because of the orthogonality of the projection effects toward the Galactic poles. These results do show us, though, that the adopted value for the Solar peculiar motion in this direction is the correct one. Figure (1) shows the effect of changing the standard $U_{\odot} = +11 \text{ km s}^{-1}$ by $\pm 3 \text{ km s}^{-1}$. An eye-ball fit from Figure (1) suggests for the U component of the solar motion a value of $U_{\odot} = +11.0 \pm 1.5 \text{ km s}^{-1}$. Also, there is no indication of a local expansion or contraction of the Galactic disk, as is also found from the kinematics of local molecular clouds (Belfort and Crovisier 1984).

A change of the orientation of the velocity ellipsoids from a cylindrical to a spherical projection tends to produce a slightly larger value for the velocity lag, especially at fainter magnitudes. The upper solid line in Fig. (2) shows that, while the fit of the model to the observed data for $V_{\odot} = +5 \text{ km s}^{-1}$, $V_{\text{LSR}}(R_0) = +270 \text{ km s}^{-1}$ is good at bright magnitudes ($B_J < 14$), the model underestimates the lag at fainter magnitudes. However, by changing the orientation of the velocity ellipsoid we can actually increase the lag. This is shown in Figure (2) where a model with $V_{\odot} = +5 \text{ km s}^{-1}$, $V_{\text{LSR}}(R_0) = +270 \text{ km s}^{-1}$, and $q = 1$ gives a better overall fit to the observed median motion in the V-component than a model with $q = 0$.

4.2. Implications for the mass of the Galaxy

The large LSR speed suggested by the SPM data poses an interesting problem to dynamical models of the Galaxy. Since the LSR is, by definition, in centrifugal equilibrium with the Galactic potential, a larger LSR velocity implies a larger mass for the Galaxy interior to the Solar circle. As emphasized by Rohlfs and Kreitschmann (1988), practically all Galactic-mass models follow, to within 10%, the relation:

$$\frac{M(r)}{10^{10} M_{\odot}} = 2.3 \times 10^{-5} \left(\frac{r}{\text{kpc}} \right) \left(\frac{V_{\text{rot}}(r)}{\text{km s}^{-1}} \right)^2 \quad (1)$$

where $M(r)$ is the mass interior to the radius r , and $V_{\text{rot}}(r)$ is the circular speed at a distance r from the Galactic center. We see that an increase in the LSR speed from 220 km s^{-1} to 270 km s^{-1} would imply as much as 50% more mass interior to the Solar circle. Is there any other evidence that this might be plausible?

An indication that this is perhaps not only possible but, indeed, dynamically required, comes from a recent measurement of the absolute proper-motions of the Large Magellanic Clouds (LMC) by Anguita et al. (1999). They have used QSOs as an inertial reference frame to obtain absolute proper-motions of stars in three fields toward the LMC. The proper-motions are based on CCD frames taken at 7 to 12 epochs spanning eight years, with a mean accuracy for the proper-motions from these three fields of $\pm 0.2 \text{ mas yr}^{-1}$. Their derived motion for the LMC is larger than that inferred by previous photographic studies that have used either bright stars or extended galaxies as a reference frame. Anguita et al. derive a mass for the Milky Way out to 50 kpc of $3-4 \times 10^{12} M_{\odot}$, in contrast to previous determinations (e.g., Lin and Lynden-Bell 1982) that implied a value of $7 \times 10^{11} M_{\odot}$. Anguita's result is fully consistent with our findings: Galactic mass models predict that the mass interior to the Solar circle is only about 4% of the total Galactic mass (Rohlfs and Kreitschmann 1988). The range of mass values found by Anguita et al. implies, through Eq. (1) for $r = 8.5 \text{ kpc}$, a rotational velocity at the Solar Galactocentric distance in the range $250 - 286 \text{ km s}^{-1}$, which indeed brackets our results. Interestingly enough, a mass of $7 \times 10^{11} M_{\odot}$ implies a very improbable rotational speed of 120 km s^{-1} .

On the other hand, our derived value for $V_{\text{LSR}}(R_0)$

seems to be in conflict with a recent determination of the proper motion of Sgr A* by Reid et al. (1999), and Backer & Sramek (1999) who find $V_{\text{LSR}}(R_0)/R_0 \approx 29 \text{ km s}^{-1} \text{ kpc}^{-1}$ (employing the good assumption that Sgr A* is at rest), which would require R_0 larger than 9 kpc, inconsistent with almost all previous determinations. Could this inconsistency point, perhaps, to evidence against our basic model assumptions which would then lead to a biased $V_{\text{LSR}}(R_0)$?

4.3. Other evidence for a larger Galactic mass

Recently, Zaritsky (1999) has reviewed most of the evidence regarding the mass of the Galactic halo and, by extension, the total mass of the Galaxy. He points out that a squared ratio of rotational circular velocities of 1.5 (such as that found from the 220 to 270 km s^{-1}) is not considered a serious discrepancy in the modeling of the total Galactic mass. He furthermore points out that timing arguments involving the relative motion of M 31 and the Milky Way imply a mass for the latter of $\sim 1.4 \times 10^{12} M_{\odot}$. A similar argument, based on the Leo I-Milky-Way system, leads to a mass of $1.1 - 1.5 \times 10^{12} M_{\odot}$. In both cases, the analysis excludes angular momentum, overlapping mass distributions at earlier times, and the evolution (growth) of the Galactic halo with age, all of which would lead to an increased total mass. Zaritsky also reviews an extension of the two-body calculations into the larger environment, which leads to an even larger mass, in the range $1.9 - 2.3 \times 10^{12} M_{\odot}$, concluding that all this evidence implies a mass out to $\sim 200 \text{ kpc}$ of $\geq 1.2 \times 10^{12} M_{\odot}$. Thus, Anguita's results based on the LMC proper motions, and its correspondence to our findings, are in line with Zaritsky's arguments.

5. Conclusions

We have used the distribution of absolute proper motions of $\sim 30,000$ randomly selected stars, part of the Southern Proper Motion survey, to constrain a structural & kinematic model of the Galaxy. The absolute proper-motions in the U-component indicate a solar peculiar motion of $11.0 \pm 1.5 \text{ km s}^{-1}$, with no need for a local expansion or contraction term. In the V-component, the absolute proper-motions can only be satisfactorily reproduced by the model if we adopt a solar peculiar motion of $+5 \text{ km s}^{-1}$, a large LSR speed of 270 km s^{-1} , and a (disk) velocity ellipsoid that always points towards the Galactic center.

The larger than expected LSR speed leads to a

larger mass for the Galaxy interior to the Solar Circle by a factor of about 1.5. This is compared with other recent evidence indicating that the Milky-Way might indeed be more massive than previously thought. This evidence comes mainly from a new measurement of the proper motion of the LMC (Anguita et al. 1999) and from the binary motion ("timing argument") of the M31 - Milky Way and Leo I - Milky Way pairs (Zaritsky 1999). Our larger LSR speed is also coincident with a similar finding by Miyamoto and Zhu (1998) using O-B5 stars and Cepheids from the Hipparcos catalogue. We stress that our result of a large LSR speed rests crucially on the assumptions of both axisymmetry and equilibrium dynamics. Failure to meet these conditions would lead to inconsistencies, such as those related to the Solar-galactocentric distance indicated at the end of Section 4.2.

High accuracy proper motion measurements for the SMC, and proper motions for some of the nearby dwarf Spheroidals will help in confirming a larger value for the mass of the Galaxy, while future space interferometric space missions like FAME (<http://www.usno.navy.mil/fame>), GAIA (<http://astro.estec.esa.nl/GAIA/>), and SIM (<http://sim.jpl.nasa.gov/>), will directly measure the disk rotational speed across the entire Galaxy.

The SPM is a joint project of the Universidad Nacional de San Juan, Argentina and the Yale Southern Observatory. Financial support for the SPM has been provided by the US NSF and the UNSJ through its Observatorio Astronómico "Félix Aguilar". We would like to also acknowledge the invaluable assistance of Lic. Carlos E. López, who participated in, and supervised, all of the SPM second-epoch observations. We are indebted to an anonymous referee who provided many useful comments on the limitations and potentialities of the Galactic model employed here. These, and other comments by the referee, have greatly helped to clarify many points of the original manuscript.

REFERENCES

- Anguita, C., Loyola, P., Pedreros, M.H., 1999, Submitted to AJ
- Backer, D.C. & Sramek, R.A., 1999, astro-ph/9906048
- Belfort, P., and Crovisier, J., 1984, A&A, 136, 368
- Delhaye, J., 1965, in Galactic Structure, Stars and Stellar Systems, Vol. 5, eds. A. Blaauw and M. Schmidt (The University of Chicago, Chicago), page 61
- Dehnen, W., Binney, J. J., 1998, MNRAS, 298, 387
- Dehnen, W., 1998, AJ, 115, 2384
- Gilmore, G., Wyse, R. F. G., and Kuijken, K., 1989, ARA&A, 27, 555
- Girard, T.M., Platais, I., Kozhurina-Platais, V., van Altena, W.F., & López, C.E. 1998, AJ, 115, 855 (Paper I)
- Kerr, F. J., and Lynden-Bell, D., 1986, MNRAS, 221, 1023
- Kuijken, K., and Gilmore, G., 1989, MNRAS, 239, 571
- Lin, D.N.C., and Lynden-Bell, D., 1982, MNRAS, 198, 707
- Méndez, R.A., & van Altena, W.F. 1996, AJ, 112, 655
- Méndez, R.A., Platais, I., Girard, T.M., Kozhurina-Platais, V., van Altena, W.F. 1999, AJ, submitted (Paper III)
- Mihalas, D., and Binney, J., 1981, Galactic Astronomy, San Francisco: Freeman
- Miyamoto, M., and Zhu, Z., 1998, AJ, 115, 1483
- Platais, I., Girard, T.M., Kozhurina-Platais, V., van Altena, W. F., López, C. E., Méndez, R. A., Ma W.-Z., Yang, T.-G., MacGillivray, H. T., Yentis, D. J., 1998, AJ, 116, 2556 (Paper II)
- Reid, M.J., Readhead, A.C.S., Vermeulen, R.C., Treuhaft, R.N., 1999, astro-ph/990575
- Robin, A. C., and Oblak, E., 1987, Evolution of Galaxies, 10th IAU European Regional Meeting, Vol. 4, J. Palous, Ondrejov: Astronomical Institute of the Czechoslovak Academy of Sciences, 323
- Rohlf, K., and Kreitschman, J., 1988, A&A, 201, 51
- Zaritsky, D. 1999, in The Third Stromlo Symposium: The Galactic halo, ASP Conf. Ser., Vol. 165, eds. B. K. Gibson, T. S. Axelrod and M.E. Putnam (Astron. Soc. of the Pacific, California), page 34

Fig. 1.— Median observed (filled dots with error bars) and predicted (lines) absolute proper-motions along U (Galactocentric direction). In some cases the error bars are smaller than the size of the dot. The solid line is for the standard model, the dashed line is for a model with a Solar peculiar motion in the U-component of $+8 \text{ km s}^{-1}$, while the dot-dashed line is for a model with $U_{\odot} = +14 \text{ km s}^{-1}$. The standard model, with $U_{\odot} = +11 \text{ km s}^{-1}$ provides the best fit to the U-component of the observed motion. The agreement to the model also indicates the absence of any significant expansion/contraction in the local disk (see text).

Fig. 2.— Median observed (filled dots with error bars) and predicted (lines) absolute proper-motions along V (Galactic rotation). In some cases the error bars are smaller than the size of the dot. The lower solid line is from the standard model with $V_{\odot} = +14 \text{ km s}^{-1}$, and $V_{\text{LSR}}(R_0) = +220 \text{ km s}^{-1}$. The dashed line is for a model with a disk having a rotational speed of $+270 \text{ km s}^{-1}$. The dotted line indicates the predictions for a model with a Solar peculiar motion in the V-component of $+5 \text{ km s}^{-1}$ instead of the classical value $+14 \text{ km s}^{-1}$ adopted in the standard model, while the triple-dot dash line shows a model with, both, $V_{\odot} = +5 \text{ km s}^{-1}$, and $V_{\text{LSR}}(R_0) = +270 \text{ km s}^{-1}$. Clearly, a larger LSR speed improves the model predictions at the fainter ($B_J \geq 15$) bins. Finally, the upper solid line is for a model with $V_{\odot} = +5 \text{ km s}^{-1}$, $V_{\text{LSR}}(R_0) = +270 \text{ km s}^{-1}$, and $q = 1$, indicating that the best fit to the observed median motion in the V-component is provided by a velocity ellipsoid whose major axis points to the Galactic center as we move away from the Galactic plane.







